

# Concentrations of persistent organochlorine contaminants in bowhead whale tissues and other biota from northern Alaska: Implications for human exposure from a subsistence diet<sup>☆</sup>

P.F. Hoekstra<sup>a,b</sup>, T.M. O'Hara<sup>c,\*</sup>, S.M. Backus<sup>a</sup>, C. Hanns<sup>c</sup>, D.C.G. Muir<sup>a</sup>

<sup>a</sup>National Water Research Institute, Environment Canada, Burlington, Ont., Canada L7R4A6

<sup>b</sup>Syngenta Crop Protection Canada, Guelph, Ont., Canada N1G4Z3

<sup>c</sup>Department of Wildlife Management, North Slope Borough, Barrow, AK 99723, USA

Received 17 June 2004; received in revised form 16 September 2004; accepted 24 September 2004

## Abstract

Bowhead whale (*Balaena mysticetus*;  $n = 5$ ) blubber, liver, muscle, kidney, heart, diaphragm, tongue, and uncooked maktak (bowhead whale epidermis and blubber) were collected during subsistence hunts at Barrow, AK, USA (1997–1999) to measure concentrations of persistent organochlorine contaminants (OCs). The exposure of humans to OCs via bowhead whales and other biota [fish, ringed (*Phoca hispida*) and bearded seals (*Erignathus barbatus*), and beluga whale (*Delphinapterus leucas*)] as part of a subsistence diet was evaluated. Concentrations of OCs in bowhead whale tissues were correlated with lipid content ( $P < 0.001$ ) and were less than levels in other marine mammals reported herein, reflecting the lower trophic status of this cetacean. The relative proportions of hexachlorobenzene (HCB) and sum ( $\Sigma$ ) concentrations of chlordane components ( $\Sigma$ CHL), DDT-related compounds ( $\Sigma$ DDT), and polychlorinated biphenyls ( $\Sigma$ PCB) were not statistically different among the tissues analyzed ( $P < 0.05$ ). However, relatively higher proportions of hexachlorocyclohexane isomers ( $\Sigma$ HCH), particularly  $\beta$ -HCH, were observed in bowhead whale heart and diaphragm ( $P < 0.03$ ). Based on Canadian and World Health Organization daily intake guidelines, “safe” human consumption rates of bowhead whale tissue and other marine biota were calculated. The most restrictive limits (mean value) for daily consumption for bowhead and beluga whale were 302 and 78 g for maktak and maktaaq (beluga whale epidermis and blubber), respectively. The tolerable daily intake limits of dioxin-like compounds from the consumption of bowhead whale blubber and liver were calculated to be 199 g (approximately 600 g for maktak) and 2222 g, respectively. A detailed profile of traditional/country foods consumed by subsistence communities of northern Alaska is required to address chronic exposure in more detail. Overall, bowhead whale tissues and other biota from northern Alaska are safe to consume at, or below, the levels calculated.

© 2004 Elsevier Inc. All rights reserved.

**Keywords:** Arctic; Cetacean; Marine mammal; Subsistence; Traditional food; Tissue distribution

<sup>☆</sup>The Department of Wildlife Management (North Slope Borough, Barrow, AK, USA) provided logistical and financial support for this study mostly via the US Department of Commerce National Oceanic and Atmospheric Administration Coastal Impact Assistance Award (No. NA170Z2054). Whale samples were provided by the Native hunters, whaling captains, and crews (Barrow Whaling Captains Association, Barrow, AK), cooperation of the Alaskan Eskimo Whaling Commission (Barrow, AK), and beluga whale hunters of Point Lay, AK. T. Rowles (Marine Mammal Health and Stranding Response Program, Washington, DC, USA), and S. Moore and M. Goshko (NOAA, Seattle, WA, USA) provided assistance in acquiring the necessary permits. Samples were transported to the National Water Research Institute (Environment Canada, Burlington, Ont., Canada) under US Export and Canadian Import permits in accordance with the Convention on International Trade in Endangered Species (US694250 and CA-CW-IM-0053-00, respectively) and via provision of the US Marine Mammal Protection Act (Permit No. 782-1399).

\*Corresponding author. Institute of Arctic Biology, University of Alaska Fairbanks, P.O. Box 757000 Fairbanks, AK 99775-7000, USA. Fax: +1 907 474 2799.

E-mail address: [fftmo@uaf.edu](mailto:fftmo@uaf.edu) (T.M. O'Hara).

## 1. Introduction

The presence of persistent organochlorine contaminants (OCs), a structurally diverse group of agricultural and industrial compounds (or byproducts), in the Arctic is well known. These chemicals were first detected in the region in the late 1960s (Jensen, 1972) and have been found in virtually every compartment in the Arctic due to their environmental recalcitrance and long-range transport from industrial and agricultural activities via atmospheric and oceanic currents (de Wit et al., 2004). The persistence, toxicity, and bioaccumulation potential of OCs is particularly significant in the arctic marine environment, where many species feed on prey with greater lipid content compared to phylogenetically similar species inhabiting lower latitude ecosystems, resulting in the accumulation of OCs to relatively high concentrations in top predatory marine species such as polar bears (*Ursus maritimus*) and ringed seals (*Phoca hispida*) (de Wit et al., 2004). The accumulation of OCs in arctic biota is of concern to indigenous peoples of this region, who may be exposed to OCs at greater concentrations than populations in southern Canada or the USA due to the consumption of lipid-rich traditional foods (Sandau et al., 2000).

The subsistence diet of indigenous peoples of the Arctic varies by region due to prey availability and local hunting practices. In northern Alaska, Inuit have hunted and consumed arctic wildlife and marine mammals, especially the bowhead whale (*Balaena mysticetus*), for many generations (Stoker and Krupnik, 1993). While the bowhead whale stock found in the Bering–Chukchi–Beaufort Seas region was greatly reduced by commercial whaling (Shelden and Rugh, 1996), this population is increasing at a rate of  $\approx 3\%$  per year (Raftery and Zeh, 1998) and remains a sociocultural and nutritionally important species to coastal Alaskan subsistence communities (Kassam, 2001).

Several studies have recognized the sociocultural and nutritional benefit of traditional foods to arctic peoples (e.g., Van Oostdam et al., 2003). However, the accumulation of OCs in Native populations from subsistence arctic communities has raised some questions concerning the suitability of terrestrial and marine wildlife from this region for human consumption (Kuhnlein and Chan, 2000; Rubin et al., 2001; Sandau et al., 2000; Van Oostdam et al., 2003). Concentrations of OCs in bowhead whale blubber and liver, along with blubber from other marine mammals from northern Alaska, have been reported (Hoekstra et al., 2002b, 2003b; Krahn et al., 2000; Kucklick et al., 2002; O'Hara et al., 1999). However, the exposure to OCs from consumption of other bowhead whale tissues as part of a subsistence diet has not been investigated.

In this paper, we discuss the potential human exposure to OCs and implications for human consump-

tion of bowhead whale tissues and other biota as part of a subsistence diet. We investigated the presence of several persistent OC groups in bowhead whale tissues that had not been examined previously (muscle, kidney, heart, diaphragm, and tongue) and uncooked bowhead whale maktak, a traditional food item comprising bowhead whale epidermis and blubber. The concentrations and relative abundance of OCs in these bowhead whale tissues are compared to foods in general in this work and will be compared for select foods purchased locally (Barrow, AK, USA) in a separate report.

## 2. Methodology

### 2.1. Sample collection

Samples were collected from 1997 to 1999 at Barrow (71°17'N, 156°45'W), Nuiqsut, AK (70°21'N, 151°02'W), and Pt. Lay, AK (69°43'N, 163°00'W) through the North Slope Borough Department of Wildlife Management (Fig. 1) as part of a larger study to investigate the trophic ecology and transfer of persistent OCs in the arctic marine environment of northern Alaska (Hoekstra et al., 2003b). Blubber samples from ringed seals (*P. hispida*;  $n = 20$  seals), bearded seals (*Erignathus barbatus*;  $n = 20$  seals), and beluga whale (*Delphinapterus leucas*;  $n = 20$  whales;  $n = 5$  whales with epidermis) were collected from Inuit subsistence harvests. Bowhead whale tissues ( $n = 5$  whales) were obtained from the Inuit's subsistence hunt with the permission of the Alaska Eskimo Whaling Commission and Barrow Whaling Captains Association (Barrow). Several species of fish, including arctic char (*Salvelinus alpinus*;  $n = 5$  fish), pink salmon (*Oncorhynchus gorbuscha*;  $n = 7$  fish), broad whitefish (*Coregonus nasus*;  $n = 19$  fish), arctic grayling (*Thymallus arcticus*;  $n = 2$  fish), and burbot (*Lota lota*;  $n = 15$ ), were provided by Native (Inuit) subsistence fishers.




Fig. 1. Sampling sites (●) of biota and fish at Pt. Lay (beluga only), Nuiqsut (arctic grayling, burbot, and broad whitefish only), and Barrow, AK, USA (1997–1999).

Field sampling techniques of mammalian tissues have been previously described (Becker et al., 1991; Hoekstra et al., 2002b). Representative samples of uncooked bowhead and beluga whale *maktak* and *maktaaq*, respectively, were prepared using an epidermis-to-blubber size ratio of 1:2 (i.e., the typically consumed dimension). Samples were initially collected in the field (from each animal) in Whirl-Pak bags and then were transported to the National Water Research Institute (Environment Canada, Burlington, Ont., Canada) under US Export and Canadian Import permits in accordance with the Convention on International Trade in Endangered Species (US694250 and CA-CW-IM-0053-00, respectively) and the US Marine Mammal Protection Act (Permit No. 782-1399). External surfaces of all samples were trimmed and removed (to minimize possible contamination from sample bags), homogenized, and stored at  $-20^{\circ}\text{C}$  in precleaned glass containers.

## 2.2. Extraction and analysis of non-planar polychlorinated biphenyls and OC pesticides

Biota samples were extracted and analyzed using previously described techniques with only minor modification (Hoekstra et al., 2002b). In brief, fish samples and full-thickness blubber cores from marine mammals were homogenized prior to OC extraction. Tissue subsamples were mixed with sodium sulfate ( $\text{Na}_2\text{SO}_4$ ) and spiked with a mixture of internal standards [two polychlorinated biphenyl (PCB) congeners, CB-30 and CB-204,  $\delta$ -HCH (hexachlorocyclohexane), 1,3,5-tribromobenzene, and 1,2,4,5-tetrabromobenzene] to monitor analyte recovery. Fish and bowhead whale tissues (except blubber and *maktak*) were weighed (10–15 g) and extracted with dichloromethane (DCM) via Soxhlet, whereas marine mammal blubber and *maktak*/*maktaaq* samples (1–3 g) were extracted with DCM using a Polytron homogenizer (Brinkmann, Westbury, NY, USA). Lipids and other bioorganic materials in each sample were removed using gel-permeation chromatography. The lipid percentage of each was determined gravimetrically on a subsample of the extract.

The analytes for each sample were concentrated and separated on 100%-activated silica gel into two fraction, hexane (Fraction 1) and hexane:DCM (1:1 by volume; Fraction 2), transferred to 2,2,4-trimethylpentane (isooctane), and concentrated. Analysis of OCs in both fractions was performed using a Hewlett-Packard 5890 gas chromatograph (Wilmington, DE, USA) with an  $^{63}\text{Ni}$ -electron capture detector (Hoekstra et al., 2002c). Sample quantification of OCs was performed using multiple external standards provided by the National Laboratory for Environmental Testing (NLET; Environment Canada, Burlington, Ont., Canada), which were analyzed after every 10 samples. The concentrations of

hexachlorobenzene (HCB) and sum ( $\Sigma$ ) group concentrations of chlordane-related products ( $\Sigma\text{CHL}$ ), DDT-compounds ( $\Sigma\text{DDT}$ ), hexachlorocyclohexane isomers ( $\Sigma\text{HCH}$ ), and  $\Sigma\text{PCB}$  are provided in Tables 1–3.

## 2.3. Coplanar PCBs and polychlorinated dibenzo-*p*-dioxin and dibenzofuran s in bowhead whale blubber and liver

A limited number of bowhead whale blubber and liver tissues ( $n = 5$ ; tissues selected from the same specimen) were randomly selected for analysis of coplanar PCBs and polychlorinated dibenzo-*p*-dioxin and dibenzofuran (PCDD/F) congeners. Other tissues were omitted from this analysis because coplanar PCB and PCDD/F concentrations in arctic wildlife are generally found at a fraction of  $\Sigma\text{PCB}$  levels (de Wit et al., 2004) and were likely to be below method detection limits.

For coplanar PCBs, bowhead whale blubber (1–2 g) and liver (5–10 g) samples were accurately weighed and homogenized with  $\text{Na}_2\text{SO}_4$ . Samples were spiked with  $^{13}\text{C}$ -labelled CB-77, CB-126, and CB-169 to monitor extraction efficiency. Samples were extracted overnight in *n*-hexane:DCM (50:50, by volume) using a Soxhlet apparatus and concentrated. Lipid percentage was determined gravimetrically using an aliquot of the extract. The remaining portion was eluted through an acidified silica gel column (22%  $\text{H}_2\text{SO}_4$  by weight, 6 g) to remove lipids and concentrated.

Coplanar PCBs were separated from non-planar OCs by high-pressure liquid chromatography (HPLC) (Bagheri et al., 1993; Lundgren et al., 2002). In brief, samples were manually injected into an Agilent 1100 HPLC system equipped with a diode-array detector (Agilent Technologies, Wilmington, DE, USA). The HPLC separation of planar from non-planar PCBs was performed on two 15 cm  $\times$  4.6 mm (internal diameter, i.d.) Cosmosil PYE-5 columns with a particle size of 5  $\mu\text{m}$  (Nacalai Tesque, Japan). Columns were connected in series and temperature was maintained at  $20^{\circ}\text{C}$ . Hexane was used for the elution of PCBs at a flow rate of  $1\text{ mL min}^{-1}$ . The fraction containing planar PCBs was collected in a quantity of between 11 and 20 mL.

The coplanar PCB fraction was fortified in isooctane and concentrated to 100  $\mu\text{L}$ . Analysis of coplanar PCBs was performed using an Agilent 6890N gas chromatograph equipped with a 5973N mass selective detector using negative chemical ionization (Bagheri et al., 1993). The gas chromatograph capillary fused-silica column used was an Agilent HP-5MS (30 m  $\times$  0.25 mm i.d.  $\times$  0.25- $\mu\text{m}$  film thickness). Helium was selected as the carrier and methane was used as the chemical reagent gas.

Extraction and analysis of PCDD/Fs in bowhead whale blubber and liver samples by Axys Analytics (Sidney, BC, Canada) used previously

Table 1

Mean ( $\pm 1$  SD), median, and range of lipid content (%) and sum ( $\Sigma$ ) concentrations of persistent organochlorine contaminant classes and HCB ( $\text{ng g}^{-1}$ , wet wt) in tissue and uncooked maktak from five ( $n=5$ ) bowhead whales (*Balaena mysticetus*) collected in 1997–1999

Tissue	% Lipid	HCB	$\Sigma\text{CHL}$	$\Sigma\text{DDT}$	$\Sigma\text{HCH}$	$\Sigma\text{PCB}$
Blubber	83.7 $\pm$ 11.0	100 $\pm$ 24	255 $\pm$ 122	377 $\pm$ 343	297 $\pm$ 126	354 $\pm$ 211
	83.0	112	239	258	273	313
	62.3–99.3	73.0–273	94.5–701	118–1925	92.2–763	153–1305
Maktak	56.1 $\pm$ 5.0	30.5 $\pm$ 5.24	95.3 $\pm$ 27.1	57.3 $\pm$ 11.6	65.7 $\pm$ 16.9	105 $\pm$ 28.2
	59.5	28.1	90.6	54.9	59.6	93.1
	49.5–60.0	26.0–39.1	69.4–136	48.1–77.1	50.1–87.4	81.8–153
Kidney	8.00 $\pm$ 5.76	5.50 $\pm$ 3.06	7.82 $\pm$ 4.72	6.37 $\pm$ 4.20	7.20 $\pm$ 5.37	12.0 $\pm$ 8.91
	7.11	5.70	7.90	5.98	6.21	10.6
	2.75–15.0	2.49–8.13	3.10–12.4	2.24–11.4	2.05–14.3	3.97–22.7
Liver	6.58 $\pm$ 1.70	3.17 $\pm$ 1.77	5.48 $\pm$ 2.22	3.72 $\pm$ 1.54	9.45 $\pm$ 2.22	9.10 $\pm$ 4.13
	6.89	3.66	5.70	3.69	9.14	9.04
	3.36–10.4	1.73–9.35	2.30–10.6	1.35–6.72	6.25–13.82	1.66–16.5
Muscle	2.39 $\pm$ 1.85	1.60 $\pm$ 0.96	2.32 $\pm$ 1.17	1.71 $\pm$ 1.01	2.74 $\pm$ 1.68	1.87 $\pm$ 0.90
	1.96	1.16	1.63	2.15	1.86	1.30
	0.80–5.34	0.69–3.02	1.20–3.69	0.59–2.81	1.41–5.48	1.18–3.19
Heart	1.65 $\pm$ 0.25	1.31 $\pm$ 0.51	1.53 $\pm$ 0.92	0.25 $\pm$ 0.18	3.50 $\pm$ 0.99	1.95 $\pm$ 0.85
	1.52	1.14	1.02	0.26	3.14	1.66
	1.45–2.05	0.98–2.21	0.99–3.13	0.03–0.46	2.39–4.88	1.26–3.38
Diaphragm	2.39 $\pm$ 2.19	1.96 $\pm$ 1.26	2.69 $\pm$ 2.06	1.59 $\pm$ 1.04	4.47 $\pm$ 2.51	3.43 $\pm$ 1.94
	0.87	1.33	1.56	0.96	3.06	2.62
	0.66–4.98	0.90–3.82	1.03–5.46	0.74–2.93	2.38–7.26	1.57–6.28
Tongue	10.2 $\pm$ 2.3	8.58 $\pm$ 1.54	18.4 $\pm$ 3.76	15.8 $\pm$ 4.67	22.0 $\pm$ 3.76	15.7 $\pm$ 1.57
	10.7	7.93	19.7	17.2	23.3	15.8
	7.11–12.4	7.60–10.9	12.9–21.4	9.13–19.8	16.5–24.9	14.1–17.3

Values provided for each parameter are arithmetic mean  $\pm 1$  SD, median, and data range. HCB, hexachlorobenzene;  $\Sigma\text{CHL}$ , sum concentration of *cis*- and *trans*-chlordanes, *cis*- and *trans*-non-achlor, and oxychlordanes;  $\Sigma\text{DDT}$  = sum of *o,p'*- and *p,p'*-DDT, DDE, and DDD;  $\Sigma\text{HCH}$ , sum of  $\alpha$ -,  $\beta$ -, and  $\gamma$ -HCH;  $\Sigma\text{PCB}$ , sum of congeners CB-4/10, 5/8, 6, 7/9, 12/13, 15/17, 16, 18, 19, 21/33/53, 22, 24/27, 25, 26, 28, 29/54, 31, 32, 40, 41/71, 42, 43, 44, 45, 46, 47/48, 49, 50, 51, 52, 55, 56/60, 59, 63, 64, 66/95, 70, 74, 76/98, 81, 82, 83, 85, 87, 91, 92/84, 97, 99, 100, 101, 105, 107/147, 110, 114, 118, 119, 128, 129/178, 130/176, 132, 133/149, 135, 136, 137, 138, 141/179, 143, 144, 151, 153, 156, 158, 163, 167, 170/190, 171, 172, 173/202, 174, 175, 177, 180, 182, 183, 185, 187, 189, 191, 193, 194, 195, 196/203, 197, 198, 199, 201, 205, 206, 207, 208, and CB-209 (IUPAC designations).

described techniques with only minor modification (US EPA, 1994). Bowhead blubber and liver samples were homogenized with  $\text{Na}_2\text{SO}_4$  and spiked with a mixture of  $^{13}\text{C}$ -labelled PCDD and PCDF isomer (as listed in Table 3) surrogate standards to monitor extraction efficiency. Samples were extracted in toluene via Soxhlet under clean-room laboratory conditions (positive pressure, carbon and HEPA filters) and concentrated. Extracts were cleaned up and PCDD/Fs were isolated through a series of chromatographic columns (US EPA, 1994). Analysis was performed using a high-resolution gas chromatograph coupled to a high-resolution mass spectrometer. Compound separation was accomplished using a DB-5 capillary chromatography column (60 m  $\times$  0.25 mm i.d.  $\times$  0.1- $\mu\text{m}$  film thickness; J&W Scientific, Folsom, CA, USA). A second column, DB-225 (30 m  $\times$  0.25 mm i.d.  $\times$  0.15- $\mu\text{m}$  film thickness), was used for confirmation of 2,3,7,8-tetraCDF.

#### 2.4. Analytical quality assurance

Recovery of non-planar PCB and OC surrogate standards ranged from 70% to 98% and concentrations

were adjusted accordingly. Detection limits ranged from 0.02 to 0.1  $\text{ng g}^{-1}$  for individual PCB congeners (including non-ortho Cl-substituted PCBs) and OC pesticides and 0.2–5.0  $\text{pg g}^{-1}$  for PCDD/F analysis. Quality assurance protocol included extraction and analysis of laboratory blanks with every batch of 10 samples and the use of two standard reference materials (SRM1588 Organics in Cod Liver Oil and SRM1945 Organics in Whale Blubber Homogenate) from the National Institute of Standards and Technology (Gaithersburg, MD, USA). Results of SRM analysis were within 20% of certified values. The laboratory that conducted coplanar PCB analyses (NLET) had successfully participated in an international interlaboratory comparison program on PCB analysis (QUASIMEME, Aberdeen, UK).

#### 2.5. Statistical analysis and calculations

Statistical analyses were conducted using the SY-STAT statistical package, Version 8.0 (SPSS, Chicago, IL, USA). All statistical tests were two-tailed and maximum probability of a Type-I error ( $\alpha$ ) was

Table 2

Mean ( $\pm 1$  SD), median, and range of lipid content (%) and sum ( $\Sigma$ ) concentrations of persistent organochlorine contaminant classes and HCB ( $\text{ng g}^{-1}$ , wet wt) in various marine mammals and fish from Alaska (1997–1999)

Species	Tissue	% Lipid	HCB	$\Sigma\text{CHL}$	$\Sigma\text{DDT}$	$\Sigma\text{HCH}$	$\Sigma\text{PCB}$
Beluga whale ( <i>Delphinapterus leucas</i> )	Blubber <i>n</i> = 20	84.8 $\pm$ 8.15	239 $\pm$ 91.4	816 $\pm$ 84.0	1979 $\pm$ 954	212 $\pm$ 161	2730 $\pm$ 1024
		84.0	233	792	2120	175	2680
		71.4–100	55.8–487	135–1936	254–4210	86.7–865	620–5248
Maktaaq <i>n</i> = 5	Blubber	44.2 $\pm$ 9.60	76.3 $\pm$ 21.5	520 $\pm$ 226	324 $\pm$ 95.3	66.7 $\pm$ 23.1	962 $\pm$ 264
		48.3	77.9	453	305	64.9	898
		28.3–53.3	39.2–106	258–922	194–483	43.3–97.0	619–1390
Ringed seal ( <i>Phoca hispida</i> )	Blubber <i>n</i> = 20	82.2 $\pm$ 13.6	17.3 $\pm$ 11.7	488 $\pm$ 400	269 $\pm$ 198	203 $\pm$ 127	691 $\pm$ 526
		86.1	12.3	400	230	141	495
		54.7–100	4.95–49.5	68.8–1484	28.6–818	37.2–1068	124–2114
Bearded seal ( <i>Erignathus barbatus</i> )	Blubber <i>n</i> = 7	79.5 $\pm$ 3.96	6.65 $\pm$ 0.55	188 $\pm$ 19.4	156 $\pm$ 21.2	89.3 $\pm$ 36.4	293 $\pm$ 30.9
		84.4	7.52	217	138	64.8	270
		60.7–90.0	3.92–8.32	123–245	111–267	25.1–300	196–411
Pink salmon ( <i>Oncorhynchus gorbuscha</i> )	Fillet <i>n</i> = 7	6.34 $\pm$ 0.26	1.45 $\pm$ 0.52	1.32 $\pm$ 0.27	1.83 $\pm$ 0.27	1.36 $\pm$ 0.58	2.63 $\pm$ 3.50
		6.46	2.10	1.18	1.63	1.58	2.89
		5.21–7.10	1.10–3.25	0.54–2.05	0.98–2.11	0.92–2.15	0.57–5.86
Arctic char ( <i>Salvelinus alpinus</i> )	Fillet <i>n</i> = 5	4.54 $\pm$ 1.79	1.44 $\pm$ 0.53	1.35 $\pm$ 1.10	1.62 $\pm$ 0.38	1.81 $\pm$ 0.88	2.34 $\pm$ 0.95
		3.64	1.23	1.29	1.48	1.65	2.54
		2.36–7.32	0.63–1.81	0.18–2.17	1.18–2.15	1.06–2.25	0.89–3.15
Broad whitefish ( <i>Coregonus nasus</i> )	Fillet <i>n</i> = 19	4.19 $\pm$ 1.04	0.47 $\pm$ 0.18	0.49 $\pm$ 0.46	0.14 $\pm$ 0.28	0.22 $\pm$ 0.09	2.89 $\pm$ 2.84
		3.98	0.42	0.39	0.06	0.22	1.64
		2.60–6.01	0.25–0.91	0.10–2.21	0.02–1.20	0.06–0.40	0.16–10.5
Arctic grayling ( <i>Thymallus arcticus</i> )	Fillet <i>n</i> = 2	4.45 $\pm$ 2.70	4.83 $\pm$ 2.59	1.28 $\pm$ 0.90	1.32 $\pm$ 0.72	0.26 $\pm$ 0.29	2.13 $\pm$ 1.71
		—	—	—	—	—	—
		2.54, 6.36	3.00, 6.67	0.64, 1.91	0.81, 1.83	0.06, 0.47	0.92, 3.34
Burbot ( <i>Lota lota</i> )	Liver <i>n</i> = 15	40.6 $\pm$ 8.93	9.74 $\pm$ 2.70	25.2 $\pm$ 14.7	16.2 $\pm$ 5.89	4.41 $\pm$ 1.58	47.3 $\pm$ 17.7
		38.9	9.96	22.7	15.8	4.55	47.2
		31.4–62.8	3.31–13.8	3.74–63.4	3.88–27.4	1.60–6.79	15.7–81.9

Values provided for each parameter are arithmetic mean  $\pm 1$  SD, median (if applicable), and data range; *n*, number of specimens analyzed.

Table 3

The mean ( $\pm 1$  SD) tolerable daily intake limits (Eq. (2)) for various subsistence dietary items based on dietary exposure guidelines<sup>a,b</sup> and the wet weight concentrations of persistent organochlorine contaminants listed in Tables 1 and 2

Species	Tissue	Tolerable daily intake limits <sup>c</sup> (g)				
		HCB	$\Sigma\text{CHL}$	$\Sigma\text{DDT}$	$\Sigma\text{HCH}$	$\Sigma\text{PCB}$
Bowhead whale	Blubber	184 $\pm$ 80	149 $\pm$ 43	4680 $\pm$ 1940	67 $\pm$ 15	218 $\pm$ 72
	Maktak	687 $\pm$ 99	389 $\pm$ 100	25,000 $\pm$ 4,270	302 $\pm$ 73	697 $\pm$ 152
	Tongue	2390 $\pm$ 352	1974 $\pm$ 494	96,800 $\pm$ 38,100	980 $\pm$ 197	4,480 $\pm$ 448
	Liver	2930 $\pm$ 1563	7594 $\pm$ 2823	> 500,000	2185 $\pm$ 411	7200 $\pm$ 1877
	Kidney	5130 $\pm$ 3050	6270 $\pm$ 4,078	> 325,000	4800 $\pm$ 3890	9510 $\pm$ 6950
	Muscle	16,200 $\pm$ 8600	18,453 $\pm$ 8476	> 500,000	9730 $\pm$ 4470	43,800 $\pm$ 17,100
	Diaphragm Heart	21,600 $\pm$ 14,400 25,300 $\pm$ 8050	20,663 $\pm$ 12,650 27,800 $\pm$ 11,935	> 500,000 > 500,000	5960 $\pm$ 2680 6400 $\pm$ 1310	26,100 $\pm$ 13,900 40500 $\pm$ 15,600
Beluga whale	Blubber	97 $\pm$ 63	43 $\pm$ 40	1316 $\pm$ 1,196	122 $\pm$ 44	32 $\pm$ 21
	Maktaaq	276 $\pm$ 106	80 $\pm$ 33	4727 $\pm$ 1,435	358 $\pm$ 127	78 $\pm$ 22
Ringed seal	Blubber	1924 $\pm$ 1010	182 $\pm$ 176	13,300 $\pm$ 11,700	212 $\pm$ 147	203 $\pm$ 144
Bearded seal	Blubber	3000 $\pm$ 328	200 $\pm$ 22	9,788 $\pm$ 1060	436 $\pm$ 113	254 $\pm$ 25
Fish <sup>c</sup>	Fillet	31,500 $\pm$ 20,800	90,646 $\pm$ 86,740	> 500,000	89,300 $\pm$ 87,500	64,700 $\pm$ 88,000
Burbot	Liver	2250 $\pm$ 468	1795 $\pm$ 712	80,330 $\pm$ 19,100	4940 $\pm$ 2250	1450 $\pm$ 390
TDI ( $\mu\text{g kg}^{-1} \text{ day}^{-1}$ )		0.3 <sup>a</sup>	0.5 <sup>b</sup>	20 <sup>a</sup>	0.3 <sup>a</sup>	1.0 <sup>a</sup>

<sup>a</sup>Health Canada (1996).

<sup>b</sup>US EPA (1999) and WHO (1993).

<sup>c</sup>TDIL values calculated from pooled fish data (except burbot).



established at 0.05. The relationship between PCB and OC concentrations (wet weight; wet wt) with the percentage of extractable lipids in various bowhead whale tissues was investigated using Model-II linear regression. All other statistical comparisons of OC concentrations were performed using lipid-adjusted concentrations to reduce the variability of lipid content on analyte concentrations (Hebert and Keenleyside, 1995). The influence of tissue type and lipid content on the relative proportion of  $\Sigma$ OCs among bowhead whale tissues was investigated using a general linear model (GLM):

$$\Sigma\text{OCs proportion} = \mu + \text{tissue} + \text{lipid} + (\text{tissue} \times \text{lipid}) + \varepsilon, \quad (1)$$

where  $\mu$  is a constant and  $\varepsilon$  is the error term. Preliminary analysis indicated that the first-order interaction term (tissue  $\times$  lipid) was not a significant factor according to Type-III sums-of-squares and was subsequently removed from the GLM. The Scheffé's method was selected to assess all a posteriori pairwise comparisons of group means deemed significant by the GLM.

$\Sigma$ PCB concentrations (wet wt) in all biota were compared (via the *Z* test) to threshold concentrations (wet wt) in food destined for human consumption that range from 0.2  $\mu\text{g g}^{-1}$  (meat), to 0.5  $\mu\text{g g}^{-1}$  (poultry), and to 2.0  $\mu\text{g g}^{-1}$  in fish as recommended by Health Canada (Hing, 1998). Calculations of tolerable daily consumption (or intake) limits (TDIL) of bowhead whale tissues and other marine biota as food were calculated using the following equation:

$$\text{TDIL (g)} = \frac{\text{TDI} (\mu\text{g kg}^{-1} \text{ day}^{-1}) \times \text{BW (kg)}}{\text{TC} (\mu\text{g g}^{-1}, \text{ wet})}, \quad (2)$$

where TC is the mean tissue concentration of  $\Sigma$ OC expressed in  $\mu\text{g g}^{-1}$  (wet wt) and BW is the assumed body weight of an average adult human consumer (70 kg). The established tolerable daily intake (TDI) guidelines indicate that daily intake should not exceed 20  $\mu\text{g kg}^{-1}$  BW of  $\Sigma$ DDT, 0.3 for  $\Sigma$ HCH, 1.0 for  $\Sigma$ PCB, and 0.27 for HCB (Health Canada, 1996). For  $\Sigma$ CHL, the provisional tolerable daily intake (PTDI) value of 0.5  $\mu\text{g kg}^{-1}$  of body wt was used (US EPA, 1999; WHO, 1993). Estimating exposure to neonates and children was not attempted in this manuscript, as separate studies addressing concentrations of OCs in cord blood, breast milk, etc. are underway.

The potential human exposure to PCDDs and other dioxin-like contaminants in bowhead whale blubber and liver was investigated by normalizing to concentrations to 2,3,7,8-tetraCDD equivalency (TEQ) using toxic equivalency factors provided in Van den Berg et al. (1998). Additionally, TCDD-equivalent TDIL values were calculated using the Canadian TDI value of

10  $\text{pg kg}^{-1}$  BW  $\text{day}^{-1}$  (assuming a 70-kg adult consumer) from Health Canada (1996).

### 3. Results and discussion

#### 3.1. OC concentrations in bowhead whale tissue

Concentrations of sum OC groups in bowhead whale tissues and other biota, including fish, are presented in Tables 1 and 2. The trophic transfer of OCs in this near-shore marine food web has been previously discussed (Hoekstra et al., 2003b). In summary, the varying concentrations and relative bioaccumulation of OCs in marine mammals and fish species mostly results from differences in the physical-chemical properties of OCs and the feeding strategy and biotransformation capacity of the species studied in this region (Hoekstra et al., 2003b).

$\text{Log}_{10}$ -transformed, wet wt  $\Sigma$ OC and HCB concentrations in the bowhead whale were significantly correlated with the lipid content of each tissue ( $P < 0.01$ ,  $r^2 = 0.81$ ). The influence of lipid content on OC concentrations is particularly evident when comparing contaminant levels in maktak and blubber. The lipid content of maktak was approximately <30% of the lipid percentage in corresponding blubber samples. However, the geometric mean concentrations of  $\Sigma$ OC in maktak were approximately 15–30% of mean blubber values (Table 1). While the decrease in the extractable portion of non-polar lipids from blubber to maktak is an important variable affecting OC concentrations, the likely heterogeneous stratification of OCs in blubber is also a significant factor (Aguilar et al., 1999). However, lipid-adjusted OC concentrations in biopsy samples of beluga whales from the Cook Inlet, Alaska were similar to full-core blubber samples obtained by necropsy (Krahn et al., 2003). Owing to the discrepancy between these studies, direct comparisons of OC data from biopsy (or maktak) and complete blubber-core samples should be conducted with caution.

In marine mammals, blubber is the major repository for many of the recalcitrant, lipophilic OCs (O'Shea, 1999). Subsequently, most investigations emphasize the determination of residue concentrations in this tissue to address various parameters related to organism health. Contaminant concentrations in cetaceans have been reported in other compartments, such as whole blood (or erythrocytes), liver, kidney, and muscle (e.g., Bruhn et al., 1995; Hoekstra et al., 2002b, 2003a; O'Shea, 1999). However, the detailed analysis of OCs in various cetacean tissues from the same specimen is limited compared to the amount of existing literature on contaminants in blubber.

The overall percentage of HCB,  $\Sigma$ DDT,  $\Sigma$ CHL, and  $\Sigma$ EPCBs to the total OC concentration in bowhead whale

was generally consistent among all tissues analyzed. The relative proportions of HCB (GLM,  $F_{7,31}=0.61$ ;  $P=0.74$ ),  $\Sigma$ DDT (GLM,  $F_{7,31}=0.45$ ;  $P=0.84$ ),  $\Sigma$ CHL (GLM,  $F_{7,31}=2.20$ ;  $P=0.08$ ), and  $\Sigma$ PCB (GLM,  $F_{7,31}=0.14$ ;  $P=0.91$ ) were not statistically different among bowhead whale tissues. While the limited number of tissue samples available in this study decreased the power of the statistical analysis, our findings are consistent with previous observations on the homogeneous distribution of lipid-normalized OC levels in cetacean tissues (Marsili and Focardi, 1997; Muir et al., 1999).

The relative abundance of  $\Sigma$ HCH in the bowhead whale was significantly influenced by tissue type (GLM,  $F_{7,31}=10.9$ ;  $P<0.001$ ). The proportions of  $\Sigma$ HCH of total OC concentrations in heart and diaphragm were significantly greater compared to other tissues (Fig. 2, top;  $P<0.03$ ). The greater percentage of  $\Sigma$ HCH in heart and diaphragm is due to the accumulation of  $\beta$ -HCH in these compartments (Fig. 2, bottom;  $P<0.01$ ). These results are consistent with the tissue-specific accumulation of HCH in rats, in which the accumulation of  $\beta$ -HCH in the heart was significantly greater than that of  $\alpha$ -HCH; the predominant isomer in all other tissues (Siddiqui et al., 2003). To our knowledge, this is the first report of isomer-specific accumulation of HCH in any mysticete. Preferential accumulation of isomers could have toxicological implications (Willett et al., 1998) and may therefore be useful for future hazard assessments.

### 3.2. Human dietary exposure to $\Sigma$ OCs and HCB

The  $\Sigma$ PCB concentrations in bowhead whale blubber exceeded the guideline for  $\Sigma$ PCB in meat (i.e., livestock) by approximately 80% ( $P<0.001$ ), but did not exceed those suggested for fish and poultry as recommended by Health Canada (Table 1,  $P>0.25$ ; Hing, 1998). All other bowhead whale tissues, along with all fish samples, did not exceed any of the suggested  $\Sigma$ PCB guidelines (Table 1,  $P<0.01$ ). Concentrations of  $\Sigma$ PCBs in the bearded seal blubber were statistically greater than the Health Canada  $\Sigma$ PCB threshold level for meat ( $P<0.01$ ), whereas  $\Sigma$ PCBs in ringed seals exceeded both meat and poultry guidelines ( $P<0.02$ ; both comparisons). As expected,  $\Sigma$ PCBs in beluga whale blubber exceeded all threshold levels, including those for fish (Table 2,  $P<0.004$ ), whereas beluga maktaaq  $\Sigma$ PCB concentrations only exceeded meat and poultry guidelines ( $P<0.001$ ). It should be noted that the maximum  $\Sigma$ PCB threshold levels were established for known or estimated consumption rates for the general Canadian population (Hing, 1998) and not those of indigenous peoples using traditional foods. Unfortunately, this information is not available for the subsistence communities in northern Alaska.

Tolerable daily consumption (or intake) limits for various marine mammal tissues and fish varied substantially among tissues and species (Table 3) and are likely better guidelines than those discussed above using




Fig. 2. The mean ( $\pm 1$  SD) proportion of  $\Sigma$ HCH to the total OC concentrations (top) and relative percentage of  $\alpha$ - and  $\beta$ -HCH isomers (to  $\Sigma$ HCH) among various bowhead whale tissues and maktak ( $n=5$ , all groups). Graph bars designated with the same letters were not statistically different (GLM, Scheffé test,  $P<0.05$ ).

tissue residue limits (i.e., threshold levels). The maximum allowable daily intake for bowhead whale blubber would be 67 g (or 469 g per week) over an entire human lifespan as established by  $\Sigma$ HCH concentrations. However, cetacean blubber is typically consumed with epidermis (collectively called maktak and maktaaq for bowhead and beluga whales, respectively) as part of a subsistence diet in northern Alaska. Thus, TDILs for bowhead whale maktak provides a more realistic exposure scenario to OCs than those values calculated from the consumption of blubber alone. The TDIL for maktak would be 687 g for HCB, 389 g for  $\Sigma$ CHL, 302 g for  $\Sigma$ HCH, 697 g for  $\Sigma$ PCB, and approximately 25,000 g for  $\Sigma$ DDT. The most restrictive TDIL for maktak (302 g per day, or 2114 g per week) indicates that a consumer can intake more than 2 kg (approximately 4.4 lb) of maktak every week for his or her entire life and be below the critical level of exposure for these OCs. The mean TDILs calculated for other bowhead whale tissues were significantly greater than those determined for blubber due to the relatively lower concentrations of  $\Sigma$ OCs in other compartments and thus can be safely consumed in larger amounts.

In the other biota analyzed, the most restrictive level of daily consumption was calculated to be 32 and 78 g for beluga whale blubber and maktaaq and 182 and 200 g for ringed seal and bearded seal blubber, respectively, based on  $\Sigma$ CHL or  $\Sigma$ PCB concentrations. Consequently, an adult can safely consume 546 g (i.e., 1.20 lb) of maktaaq per week, every week, for an entire lifetime. While beluga blubber is typically consumed as a part of maktaaq, seal blubber is rendered into oil and used as a dip for a variety of frozen and boiled meats and fish by Native Alaskans (Kassam, 2001); therefore, it is more difficult to provide “safe” consumption advice. The maximum TDIL was 31,500 g for HCB in the fish fillets analyzed, whereas the tolerable consumption of burbot liver was more restrictive at approximately 1450 g per day (i.e., 10,150 g or 22.4 lb per week). Given the relatively large amounts of fish and seal oil dip needed to surpass intake guidelines, these species should be considered safe to consume under the guidelines discussed.

It should be noted that the TDIL for bowhead whale tissues and other biota are subject to interpretation due to the discrepancy of  $\Sigma$ CHL TDI values between various regulatory agencies. The WHO provisional TDI value of  $0.5 \mu\text{g kg}^{-1} \text{day}^{-1}$  used in this study was calculated by applying a 100-fold “safety factor” (to account for inter- and intraspecies variability) from the chronic “no adverse effect level” (NOAEL) of  $50 \mu\text{g kg}^{-1}$  using rats exposed to technical chlordane via dietary exposure (WHO, 1993). The same TDI value (or oral reference dose) is provided by the US Environmental Protection Agency (US EPA), which is based on applying a 1/300 uncertainty factor to an

NOAEL value of  $150 \mu\text{g kg}^{-1} \text{day}^{-1}$  derived from a laboratory experiment in which mice were orally exposed to  $\Sigma$ CHL for 104 weeks (Khasawinah and Grutsch, 1989; US EPA, 1999). However, the Health Canada guideline value of  $0.05 \mu\text{g kg}^{-1} \text{day}^{-1}$  for  $\Sigma$ CHL exposure is lower and more conservative (Health Canada, 1996), reflecting a high uncertainty factor (1000) compared to other jurisdictions. However, information on how this Health Canada dietary guideline value was established is not available. Nevertheless, the application of the Health Canada TDI value for  $\Sigma$ CHL would decrease the TDIL for bowhead whale tissues by one order of magnitude (i.e., 14 and 39 g of blubber and maktak, respectively). Based on the data provided by the US EPA and WHO, it is our opinion that the Health Canada guideline is overly protective and would place undue restriction on the consumption of the subsistence foods presented in this manuscript. Therefore, the lifetime exposure guideline of  $0.05 \mu\text{g kg}^{-1} \text{day}^{-1}$  for  $\Sigma$ CHL proposed by Health Canada was not used for this analysis.

### 3.3. PCDDs and dioxin-like OCs in bowhead whale blubber and liver

Relatively low concentrations of coplanar PCBs (i.e., non-ortho substituted congeners), mono-ortho Cl-substituted PCB congeners, and PCDD and PCDF isomers were detected in bowhead whale blubber and liver samples (Table 4) compared to other OCs. Overall, the profiles of PCDDs and other dioxin-like compounds in bowhead whale blubber are consistent with reports in other cetaceans (e.g., Gauthier et al., 1998; Jarman et al., 1996; Muir et al., 1996). The compound with the largest contribution to total TEQ values is the coplanar PCB congener, 3,3',4,4',5-pentaCB (CB-126), which comprised 33 and 72% of the total TEQ value in blubber and liver, respectively. The mono-ortho Cl-substituted CB-118 and CB-156 were major contributors to the total TEQ (CB-118 blubber, 22%; liver, 6.0%; CB-156 blubber, 18%; liver, 10%) compared to other congeners. While the mono-ortho Cl-substituted PCB congeners CB-123 and CB-157 were not quantified in this study, the abundance of these congeners in arctic biota is minor (de Wit et al., 2004), and their subsequent contribution to the total TEQ is expected to be minimal.

The TCDD-TEQ concentrations were comparable to those reported in various commercially available food commodities with similar lipid content (Himberg, 1993; Dyke and Stratford, 2002; Schecter et al., 1997). For example, wet-weight TEQ concentrations of PCDD/Fs in various food items (e.g., beef, pork, chicken, farm-raised fish, etc.) ranged from 0.03 to  $2.90 \text{ pg g}^{-1}$ . However, these TEQ concentrations in the commercial foods did not include coplanar PCBs, which may



Table 4

Mean ( $\pm 1$  SD) concentration ( $\text{pg g}^{-1}$ , wet weight) and toxic equivalent (TEQ) values for polychlorinated dibenzo-*p*-dioxins (PCDDs), dibenzofurans (PCDFs), and dioxin-like polychlorinated biphenyl (PCB) congeners in bowhead whale blubber and liver ( $n = 5$ )

Compound	Concentration ( $\text{pg g}^{-1}$ wet wt)		TEF <sup>a</sup>	TEQ ( $\text{pg g}^{-1}$ wet wt)	
	Blubber	Liver		Blubber	Liver
<b>PCDDs</b>					
1,2,3,4,6,7,8-heptaCDD	ND <sup>b</sup>	$0.35 \pm 0.18$	0.01	—	$0.003 \pm 0.002$
OCDD	ND	$1.60 \pm 0.42$	0.0001	—	$0.002 \pm 0.001$
$\Sigma$ PCDDs	—	$1.95 \pm 0.39$	—	—	$0.004 \pm 0.002$
<b>PCDFs</b>					
ND	ND	ND	—	—	—
<b>Non-ortho Cl-substituted PCBs</b>					
3,3',4,4'-tetraCB (CB-77)	$8.75 \pm 4.27$	$5.50 \pm 1.73$	0.0001	$0.0009 \pm 0.0001$	$0.0006 \pm 0.0002$
3,4,4',5-tetraCB (CB-81)	$4.95 \pm 3.99$	ND	0.0001	$0.0005 \pm 0.0004$	—
3,3',4,4',5-pentaCB (CB-126)	$11.52 \pm 4.55$	$2.28 \pm 0.58$	0.1	$1.152 \pm 0.455$	$0.228 \pm 0.058$
3,3',4,4',5,5'-hexaCB (CB-169)	$9.42 \pm 4.52$	$1.49 \pm 0.73$	0.01	$0.094 \pm 0.045$	$0.015 \pm 0.007$
$\Sigma$ no-PCBs	$39.8 \pm 16.1$	$9.27 \pm 2.72$	—	$1.24 \pm 0.438$	$0.243 \pm 0.065$
<b>Mono-ortho Cl-substituted PCBs</b>					
2,3,3',4,4'-pentaCB (CB-105)	$2418 \pm 1175$	$86.5 \pm 40.9$	0.0001	$0.242 \pm 0.118$	$0.008 \pm 0.004$
2,3,4,4',5-pentaCB (CB-114)	$1200 \pm 763$	$18.5 \pm 15.3$	0.0005	$0.600 \pm 0.381$	$0.009 \pm 0.007$
2,3',4,4',5-pentaCB (CB-118)	$7870 \pm 4170$	$197 \pm 106$	0.0001	$0.787 \pm 0.417$	$0.019 \pm 0.011$
2,3,3',4,4',5-hexaCB (CB-156)	$1298 \pm 1200$	$85.2 \pm 22.6$	0.0005	$0.649 \pm 0.601$	$0.031 \pm 0.023$
2,3',4,4',5,5'-hexaCB (CB-167)	$479 \pm 393$	$7.0 \pm 5.0$	0.00001	$0.005 \pm 0.004$	$0.001 \pm 0.001$
2,3,3',4,4',5,5'-heptaCB (CB-189)	$304 \pm 230$	ND	0.0001	$0.030 \pm 0.023$	—
$\Sigma$ mo-PCBs	$13500 \pm 7420$	$376 \pm 200$	—	$2.31 \pm 1.43$	$0.071 \pm 0.010$
$\Sigma$ TEQ				$3.52 \pm 1.12$	$0.315 \pm 0.031$
$\Sigma$ TEQ range				(2.01–4.95)	(0.293–0.370)

<sup>a</sup>Toxic equivalency factors (TEFs) for PCDD/F and non- and mono-*ortho* substituted PCB congeners from Van den Berg et al. (1998).

<sup>b</sup>Not detected at concentrations greater than method detection limits.

represent a significant contribution to the overall  $\Sigma$ TEQ value (Huwe, 2002).

Based on the TDI value of  $10 \text{ pg kg}^{-1} \text{ BW day}^{-1}$  (Health Canada, 1996), and assuming a 70-kg adult consumer, the TDIL of dioxin-like compounds from the consumption of bowhead whale blubber and liver was calculated to be 199 g (range, 141–348 g) and 2222 g (range, 1892–2390 g), respectively. As previously mentioned, bowhead whale blubber is typically consumed as maktak, and, therefore, actual exposure levels to OC via consumption will be overestimated only considering blubber concentrations. By comparing differences in  $\Sigma$ OC concentrations between these two food matrices (Table 1), the TCDD–TEQ concentration in maktak is likely three times lower than blubber concentrations, thereby increasing the TDIL to approximately 600 g (i.e., 4.2 kg, or approximately 9.2 lb per week) for this food item. Consequently, the risk to human health associated with exposure to PCDDs and dioxin-like compounds is minimal considering the current toxicological information and contaminant data.

### 3.4. Limitations and uncertainty

Careful interpretation of the dietary exposure values reported in this study is required. For example, the effect of food preparation on OC concentrations was not studied. Previous investigations have noted a reduction

of  $\Sigma$ PCB and other OC concentrations in fish due to cooking (Kamrin and Fischer, 1999; Zabik et al., 1995). While the effect of rendering, fermentation, and other food preparation practices used by Alaska Natives (Kassam, 2001) on OC concentrations is unknown, the estimated daily intake values calculated from raw tissues in this study likely overestimate OC exposure from these subsistence dietary items when cooked. In addition, there are limitations in using TDI or PTDI guidelines to evaluate dietary exposure to anthropogenic compounds. As previously stated, these guidelines assume a life-long exposure and are developed using the NOAEL data from laboratory- or community-based exposure scenarios. These values were subsequently adjusted using conservative safety factors, typically by dividing the NOAEL by 100 or 1000, to compensate for interspecies differences between the test species and humans (Waltner-Toews and McEwen, 1994).

In addition, it should be emphasized that TDI/PTDI values are based on a lifetime of daily exposure. However, most wildlife and fish are hunted and consumed during select months of the year. For example, beluga whales in Wainwright, AK are generally hunted from June to July, whereas bowheads are hunted during two seasonal events in the Fall (September–October) and early Spring (April–May) at Barrow (Kassam, 2001; Fuller and George, 1997). Therefore, the estimated consumption of approximately 300 g of whale

tissue (e.g., maktak) per day over an entire lifetime is likely very conservative and overestimates the true degree of risk associated with dietary exposure to OCs from these subsistence foods. However, it should be noted that dietary surveys of these subsistence species are required to properly elucidate the extent of contaminant exposure via dietary intake. The TDIL values determined in this study only considered PCB/OC intake from each individual tissue/food item. Thus, dietary surveys are needed to consider contaminant intakes from all possible dietary sources, along with the seasonal nature of hunting practices, when evaluating which food items significantly contribute to PCB/OC exposure via an actual subsistence diet (e.g., Van Oostdam et al., 2003).

The obvious sociocultural and nutritional benefits of a subsistence diet to the residents of northern Alaska must be considered. Preliminary information on the nutritional quality of bowhead whale tissues and other wildlife suggests that these subsistence dietary items are excellent sources of nutritionally essential minerals, fatty acids, and fat-soluble vitamins, such as vitamins A, D, and E (e.g., Blanchet et al., 2000; Kenny et al., 2004; Nobmann et al., 1992; Nobmann and Lanier, 2001; Van Oostdam et al., 2003; Woshner et al., 2001). The increasing prevalence of “Western” or modern diets, instead of traditional foods, is thought to have negative health consequences, in part due to the increased consumption of total fat, saturated fats, and sucrose above recommended levels (Nobmann and Lanier, 2001; Van Oostdam et al., 2003). Upon considering the risks associated with exposure to contaminants relative to the known benefits of traditional foods, as well as the risks associated with eating poor replacement foods (e.g., Egeland et al., 1998; Kuhnlein and Chan, 2000), the consumption of bowhead whale tissues and other marine biota by Inuit of northern Alaska should be maintained and encouraged. Using a balanced approach to nutrients and contaminants, this information will allow individuals to decide for themselves which dietary habits best suit their lifestyle.

#### 4. Conclusions

Persistent organochlorine contaminants were quantified in numerous tissues of the bowhead whale and other biota. In general, concentrations were relatively low in bowhead whales compared to other marine mammals and reflect the trophic status of this cetacean (Hoekstra et al., 2002a, 2003b). Lipid content significantly influenced the concentrations of  $\Sigma$ OCs in bowhead whale tissues and accumulation profiles of  $\Sigma$ OC groups were generally homogeneous in bowhead whale tissues. However, cardiac and diaphragm tissues contained a relatively greater percentage of HCH isomers, especially

$\beta$ -HCH, compared to other tissues.  $\Sigma$ OC concentrations were used to evaluate the human consumption rates based on established exposure guidelines. Overall, bowhead whale tissues and other biota from northern Alaska are safe to consume at, or below, the levels calculated in this study. Considering the safety factors applied to establish many of the TDILs, one could exceed the suggested intake levels by 100-fold the TDIL before achieving a “no observed effect” level and thus allow for a rather large tolerable range of intake. Additional information on the dietary profile (e.g., rates of consumption, peak seasons of consumption) of traditional/country foods consumed by the subsistence communities of northern Alaska is required to more definitively address the potential human health risks resulting from chronic exposure to OCs from a subsistence-based diet and the level of nutrient intake.

#### Acknowledgments

The North Slope Borough Department of Wildlife Management (Barrow, AK, USA) initiated this study and provided logistical and financial support. Additional funding was also provided by the US Department of Commerce via the National Oceanic and Atmospheric Administration Coastal Impact Assistance Award (No. NA170Z2054). The authors gratefully acknowledge the generous provision of bowhead whale samples by the Native hunters, whaling captains, and crews (Barrow Whaling Captains Association, Barrow, AK) and the cooperation of the Alaskan Eskimo Whaling Commission (Barrow, AK) and beluga whale hunters of Point Lay, AK. Sample collection and logistics related to the shipment of samples to analytical laboratories were conducted by B. Akootchook, L. Dehn, J.C. George, C. Rosa, R. Suydam, S. Visalli, V. Woshner, and many others. We thank T. Rowles (Marine Mammal Health and Stranding Response Program, Washington, DC, USA), S. Moore, and M. Goshko (NOAA, Seattle, WA, USA) for assistance in acquiring the necessary permits. Lori Verbrugge and S. Arnold (Environmental Public Health Program, State of Alaska Section of Epidemiology, Anchorage, AK) and Craig George and Robert Suydam (Department of Wildlife Management) provided valuable reviews of an early version of the manuscript. Analytical support from Axyx Analytics (Sidney, BC, Canada) and Angela Baitz and Lisa Bradley (Environment Canada, Burlington, ON, Canada) was appreciated.

#### References

- Aguilar, A., Borrell, A., Pastor, T., 1999. Biological factors affecting variability of persistent pollutants levels in cetaceans. *J. Cetacean Res. Manage.* 1, 83–116.

- Bagheri, H., Leonards, P.E.G., Ghijsen, R.T., Brinkman, U.A.T.H., 1993. Gas chromatography-negative ion chemical ionization mass spectrometry for the determination and identification of planar polychlorinated biphenyls in biological samples. *Int. J. Environ. Anal. Chem.* 50, 257–268.
- Becker, P.R., Wise, S.A., Koster, B.J., Zeisler, R., 1991. Alaska Marine Mammal Tissue Archival Project: Revised Collection Protocol. US National Institute of Standards and Technology (NIST) Interagency Report 4529. US Dept. of Commerce, NIST, Gaithersburg, MD, USA.
- Blanchet, C., Dewailly, É., Ayotte, P., Bruneau, S., Receveur, O., Holub, B.J., 2000. Contribution of selected traditional and market foods to the diet of Nunavik Inuit women. *Can. J. Diet. Pract. Res.* 61, 50–59.
- Bruhn, R., Kannan, N., Petrick, G., Schulz-Bull, D.E., Duinker, J.C., 1995. CB pattern in the harbour porpoise: bioaccumulation, metabolism and evidence of cytochrome P450 IIB activity. *Chemosphere* 31, 3721–3732.
- de Wit, C.A., Fisk, A.T., Hobbs, K.E., Muir, D.C.G., Kallenborn, R., Krahn, M.M., Norstrom, R.J., Skaare, J., 2004. Persistent organochlorine pollutants. In: de Wit, C.A., Fisk, A.T., Hobbs, K.E., Muir, D.C.G. (Eds.), *Arctic Monitoring and Assessment Program (AMAP) II. Arctic Monitoring and Assessment Program*, Oslo.
- Dyke, P.H., Stratford, J., 2002. Changes to the TEF schemes can have significant impacts on regulation and management of PCDD/F and PCB. *Chemosphere* 47, 103–116.
- Egeland, G.M., Feyk, L.A., Middaugh, J.P., 1998. The use of traditional foods in a healthy diet in Alaska: risks in perspective. *State of Alaska Epidemiology Bulletin*, vol. 2(1) January.
- Fuller, A.S., George, J.C., 1997. Evaluation of Subsistence Harvest Data from the North Slope Borough 1993 Census for Eight North Slope Villages for the Calendar Year 1992. Department of Wildlife Management, Barrow, AK, USA.
- Gauthier, J.M., Pelletier, É., Brochu, C., Moore, S., Metcalfe, C.D., Béland, P., 1998. Environmental contaminants in tissues of a neonate St. Lawrence beluga whale (*Delphinapterus leucas*). *Mar. Pollut. Bull.* 36, 102–108.
- Health Canada, 1996. Values Presently Used by Contaminants Toxicology Section. Food Directorate, Ottawa, Ont., Canada.
- Hebert, C.E., Keenleyside, K.A., 1995. To normalize or not to normalize? Fat is the question. *Environ. Toxicol. Chem.* 14, 801–807.
- Himberg, K.K., 1993. Coplanar polychlorinated biphenyls in some Finnish food commodities. *Chemosphere* 27, 1235–1243.
- Hing, M.C., 1998. A database for environmental contaminants in traditional foods in northern and arctic Canada: developments and applications. *Food Addit. Contam.* 15, 127–134.
- Hoekstra, P.F., Dehn, L.A., George, J.C., Solomon, K.R., Muir, D.C.G., O'Hara, T.M., 2002a. Trophic ecology of bowhead whales (*Balaena mysticetus*) compared with that of other arctic marine biota as interpreted from carbon-, nitrogen-, and sulfur-isotope signatures. *Can. J. Zool.* 80, 223–231.
- Hoekstra, P.F., O'Hara, T.M., Pallant, S.J., Muir, D.C.G., 2002b. Bioaccumulation of organochlorine contaminants in bowhead whales (*Balaena mysticetus*) from Barrow, Alaska. *Arch. Environ. Contam. Toxicol.* 42, 497–507.
- Hoekstra, P.F., O'Hara, T.M., Teixeira, C., Backus, S., Fisk, A.T., Muir, D.C.G., 2002c. Spatial trends and bioaccumulation of organochlorine pollutants in marine zooplankton from the Alaskan and western Canadian Arctic. *Environ. Toxicol. Chem.* 21, 575–583.
- Hoekstra, P.F., Letcher, R.J., O'Hara, T.M., Backus, S.M., Solomon, K.R., Muir, D.C.G., 2003a. Hydroxylated and methylsulfone-containing metabolites of polychlorinated biphenyls in the plasma and blubber of bowhead whales (*Balaena mysticetus*). *Environ. Toxicol. Chem.* 22, 2650–2658.
- Hoekstra, P.F., O'Hara, T.M., Fisk, A.T., Borgå, K., Solomon, K.R., Muir, D.C.G., 2003b. Trophic transfer of persistent organochlorine contaminants (OCs) within an arctic marine food web from the southern Beaufort–Chukchi Seas. *Environ. Pollut.* 124, 509–522.
- Huwe, J.K., 2002. Dioxins in food: a modern agricultural perspective. *J. Agric. Food Chem.* 50, 1739–1750.
- Jarman, W.M., Norstrom, R.J., Muir, D.C.G., Rosenberg, B., Simon, M., Baird, R.W., 1996. Levels of organochlorine compounds, including PCDDs and PCDFs, in the blubber of cetaceans from the west coast of North America. *Mar. Pollut. Bull.* 32, 426–436.
- Jensen, S., 1972. The PCB story. *Ambio* 1, 123–131.
- Kamrin, M.A., Fischer, L.J., 1999. Current status of sport fish consumption advisories for PCBs in the Great Lakes. *Regul. Toxicol. Pharmacol.* 29, 175–181.
- Kassam, K.-A.S., 2001. Passing on the knowledge: mapping human ecology in Wainwright, Alaska. Arctic Institute of North America, University of Calgary, Calgary, AB, Canada.
- Kenny, D.E., O'Hara, T.M., Chen, T.C., Lu, Z., Tian, X., Holick, M.F., 2004. Vitamin D content in Alaskan Arctic zooplankton, fishes and marine mammals. *Zool. Biol.* 23, 33–43.
- Khasawinah, A.M., Grutsch, J.F., 1989. Chlordane: 24-month tumorigenicity and chronic toxicity test in mice. *Regul. Toxicol. Pharmacol.* 10, 244–254.
- Krahn, M.M., Burrows, D.G., Stein, J.E., Becker, P.R., Schantz, M.M., Muir, D.C.G., O'Hara, T.M., Rowles, T., 2000. White whales (*Delphinapterus leucas*) from three Alaskan stocks: concentrations and patterns of persistent organochlorine contaminants in blubber. *J. Cetacean Res. Manage.* 1, 239–249.
- Krahn, M.M., Herman, D.P., Ylitalo, G.M., Sloan, C.A., Burrows, D.G., Hobbs, R.C., Mahoney, B.A., Yanagida, G.K., Calambokidis, J., 2003. Blubber stratification in white whales and killer whales: variability in contaminant concentrations, fatty acid profiles, lipid percent and lipid class profiles that has implications for biopsy sampling. Paper SC/55/E3 presented to the International Whaling Commission (IWC) Scientific Committee, June 2003, 22pp.
- Kucklick, J.R., Struntz, W.D.J., Becker, P.R., York, G.W., O'Hara, T.M., Bohonowych, J.E., 2002. Persistent organochlorine pollutants in ringed seals and polar bears collected from northern Alaska. *Sci. Total Environ.* 287, 45–59.
- Kuhnlein, H.V., Chan, H.M., 2000. Environment and contaminants in traditional food systems of northern indigenous peoples. *Annu. Rev. Nutr.* 20, 595–626.
- Lundgren, K., van Bavel, B., Tysklind, M., 2002. Development of a high-performance liquid chromatography carbon column based method for the fractionation of dioxin-like polychlorinated biphenyls. *J. Chromatogr. A* 962, 79–93.
- Marsili, L., Focardi, S., 1997. Chlorinated hydrocarbon (HCB, DDTs and PCBs) levels in cetaceans stranded along the Italian coasts: an overview. *Environ. Monit. Assess.* 45, 129–180.
- Muir, D.C.G., Ford, C.A., Rosenberg, B., Norstrom, R.J., Simon, M., Béland, P., 1996. Persistent organochlorines in beluga whales (*Delphinapterus leucas*) from the St. Lawrence River estuary. I. Concentrations and patterns of specific PCBs, chlorinated pesticides and polychlorinated dibenzo-*p*-dioxins and dibenzofurans. *Environ. Pollut.* 93, 219–234.
- Muir, D.C.G., Braune, B., DeMarch, B., Norstrom, R., Wagemann, R., Lockhart, L., Hargrave, B., Bright, D., Addison, R., Payne, J., Reimer, K., 1999. Spatial and temporal trends and effects of contaminants in the Canadian Arctic marine ecosystem: a review. *Sci. Total Environ.* 230, 83–144.
- Nobmann, E.D., Lanier, A.P., 2001. Dietary intake among Alaska Native women resident of Anchorage, Alaska. *Int. J. Circumpolar Health* 60, 123–137.
- Nobmann, E.D., Byers, T., Lanier, A.P., Hankin, J.H., Jackson, M.Y., 1992. The diet of Alaska Native adults: 1987–1988. *Am. J. Clin. Nutr.* 1024–1032.

- O'Hara, T.M., Krahn, M.M., Boyd, D., Becker, P.R., Philo, L.M., 1999. Organochlorine contaminant levels in Eskimo harvested bowhead whales of arctic Alaska. *J. Wildl. Dis.* 35, 741–752.
- O'Shea, T.J., 1999. Environmental contaminants and marine mammals. In: Reynolds, J.E., Rommel, S.A. (Eds.), *Biology of Marine Mammals*. Smithsonian Inst. Press, Washington, DC, pp. 485–536.
- Raftery, A.E., Zeh, J.E., 1998. Estimating bowhead whale population size and rate of increase from the 1993 census. *J. Am. Stat. Assoc.* 93, 1–13.
- Rubin, C.H., Lanier, A., Socha, M., Brock, J.W., Kieszak, S., Zahm, S., 2001. Exposure to persistent organochlorines among Alaska Native women. *Int. J. Circumpolar Health* 60, 157–169.
- Sandau, C.D., Ayotte, P., Dewailly, E., Duffe, J., Norstrom, R.J., 2000. Analysis of hydroxylated metabolites of PCBs (OH-PCBs) and other chlorinated phenolic compounds in whole blood from Canadian Inuit. *Environ. Health Perspect.* 108, 611–616.
- Schecter, A., Cramer, P., Boggess, K., Stanley, J., Olson, R.J., 1997. Levels of dioxins, dibenzofurans, PCBs and DDE congeners in pooled food samples collected in 1995 at supermarkets across the United States. *Chemosphere* 34, 1437–1447.
- Shelden, K.E.W., Rugh, D.J., 1996. The bowhead whale, *Balaena mysticetus*: its historic and current status. *Mar. Fish. Rev.* 57, 1–20.
- Siddiqui, M.K.J., Nigam, U., Kaul, P.P., Seth, T.D., 2003. Bioaccumulation of HCH isomers in different tissues of young and old rats: a comparison. *Bull. Environ. Contam. Toxicol.* 56, 896–902.
- Stoker, S.W., Krupnik, I.I., 1993. Subsistence whaling. In: Burns, J.J., Montague, J.J., Cowles, C.J. (Eds.), *The Bowhead Whale*. Allen Press, Lawrence, KS, USA, pp. 579–630.
- US EPA (US Environmental Protection Agency), 1994. Method 1613: Tetra- through Octa-chlorinated Dioxins and Furans by Isotope Dilution HRGC/HRMS, Revision B. US EPA, Washington, DC.
- US EPA (US Environmental Protection Agency), 1999. Integrated Risk Information System (IRIS)—Chlordane. National Center for Environmental Assessment, Office of Research and Development, Washington, DC, <http://www.epa.gov/iris/search.htm> (accessed January 2004).
- Van den Berg, M., Birnbaum, L.S., Bosveld, A.T.C., Brunström, B., Cook, P., Feeley, M., Giesy, J.P., Hanberg, A., Hasegawa, R., Kennedy, S.W., Kubiak, T., Larsen, J.C., van Leeuwen, F.X.R., Liem, A.K.D., Nolt, C., Peterson, R.E., Poellinger, L., Safe, S.H., Schrenk, D., Tillit, D., Tysklind, M., Younes, M., Wærn, F., Zacharewski, T., 1998. Toxic equivalency factors (TEFs) for PCBs, PCDDs, PCDFs for humans and wildlife. *Environ. Health Perspect.* 106, 775–792.
- Van Oostdam, J., Donaldson, S., Feeley, M., Tremblay, N., 2003. Human health. In: *Canadian Arctic Contaminants Assessment Report II*. Ministry of Public Works and Government Services Canada, Ottawa, pp. 127.
- Waltner-Toews, D., McEwen, S.A., 1994. Chemical residues in foods of animal origin: overview and risk assessment. *Prev. Vet. Med.* 20, 161–178.
- WHO (World Health Organization), 1993. Pesticide Residues in Food—1994. Report of the joint meeting of the FAO Panel of Experts of Pesticide Residues in Food and the Environment and WHO Toxicological and Environmental Core Assessment groups. FAO Plant Production and Protection Paper 127. WHO, Geneva.
- Willett, K.L., Ulrich, E.M., Hites, R.A., 1998. Differential toxicity and environmental fates of hexachlorocyclohexane isomers. *Environ. Sci. Technol.* 32, 2197–2206.
- Woshner, V.M., O'Hara, T.M., Bratton, G.R., Suydam, R.S., Beasley, V.R., 2001. Concentrations and interactions of selected essential and non-essential elements in bowhead and beluga whales of arctic Alaska. *J. Wildl. Dis.* 37, 693–710.
- Zabik, M.E., Zabik, M.J., Booren, A.I.M., Nettles, M., Song, J.-H., Welch, R., Humphrey, H., 1995. Pesticides and total polychlorinated biphenyls in Chinook salmon and carp harvested from the Great Lakes: effects of skin-on and skin-off processing and selected cooking methods. *J. Agric. Food Chem.* 43, 993–1001.